

Guideline: Adult Central Line Guideline

Revised Date: June 2023

Review Date: June 2025

Content Experts

Patrick Duffy, MD
Anne Wagner, MD

Table of Contents

I. Population..... 2

II. General Principals 2

III. Interventions..... 2

I. Population:

The following protocol applies to burn patients in the Burn ICU.

II. General Principals

- All non-tunneled lines placed at an outside facility should be considered for removal at the earliest opportunity.
- All intravascular catheters not placed under strict sterile conditions (pre-arrival at VUMC, or emergently in the trauma bay or placed under time-critical conditions in a patient with profound physiologic instability or undergoing active cardiopulmonary resuscitation) must be removed within 48 hours.
- Antibiotic impregnated catheters should be utilized.
- Ultrasound guidance is expected for IJ placements and encouraged for subclavian and femoral placements.
- All central lines should be removed at the earliest opportunity once they are no longer clinically indicated or alternative PIV access can be established.
- When long term venous access is indicated, a Midline catheter is preferred over a PICC unless a specific need for central venous access is required for therapeutic administration.
- Central line catheter tips are not to be sent for routine culture.

III. Insertion Site Selection

- Central lines will be inserted in non-burned areas when available, preferably >5cm from the nearest burn to decrease risk of bacterial contamination/colonization.
- Femoral site is preferred if through unburned skin.
- If multiple sites are available, consider operator skill and technique in the selection process with Subclavian and IJ sites preferred over femoral placement due to increasing infection risk.

IV. Technique

- VUMC principals of insertion technique should be followed according to VUMC Policy Manual
- General components of Insertion Technique include:
 - Informed Consent
 - Time-out
 - Pain management
 - Patient positioning (Trendelenburg for lines above the diaphragm)
 - Patient monitoring with telemetry
 - Sterile Technique with full barrier precautions and PPE
 - Skin prep with CHG or Betadine if patient is undergoing Cultured Epithelial Autografting.
 - Confirmation of venous puncture through transduction or direct ultrasonographic visualization of the wire within the vein prior to dilation of the puncture.
 - Secure with 2-0 silk sutures at a minimum of three points
 - Cover with site appropriate anti-infective dressing
 - Confirm central line placement of Subclavian and IJ lines by CXR prior to infusion.
 - Exchange over wire technique using Double Gloves

- Lines being exchanged over a wire require **double sterile glove** technique for placement, where the outermost glove is to be removed following removal of the existing catheter and prior to handling the new catheter.

V. Management, Maintenance and Care

Line inserted through burned skin:

- When it is necessary to insert lines through burned skin, these lines will be changed every 8 DAYS with one exchange over a wire if no suspicion of central line infection exists followed by a new site.
- Dressing: If standard CHG dressing will not remain intact, cover insertion site with Mepilex AG cut to fit as a drain sponge dressing. Change dressing daily or PRN if soiled.

Femoral lines:

- Femoral lines are preferred if other sites would be through burned skin.
- Femoral central lines will be rotated every 8 days, once by new catheter exchange over wire then followed by a new site stick.

Subclavian and IJ lines:

- Central lines placed under sterile technique through unburned skin >5cm from the nearest burn will be evaluated every shift and maintained per VUMC policy with CHG barrier clear occlusive dressing. Routine catheter changes are not required.
- Malfunctioning lines are eligible for exchange over wire under double glove sterile technique.
- Lines suspected of infection should be exchanged for a new catheter at a new site. If the new site is contraindicated, exchange over wire may be performed.
- Evidence of local catheter infection such as purulence, erythema, or tenderness, mandates a catheter change to a new site.

Lines suspected of infection:

- When there are signs of active infection around the central line insertion site, the line should be removed as soon as possible.
- Lines suspected of infection are not eligible for exchange over a wire.
- Patients with evidence of bloodstream infection and SIRS/Sepsis require a CVC line change to a new site.

VI. Special considerations

Patients with Deep Venous Thrombosis

- Consider Interventional Radiology consultation for placement if the access site would pass through the clot burden.

Patients on anticoagulants

- Select a compressible site for insertion when available.
- Utilize micro-puncture technique for access.

VII. References

- VUMC Policy Manual. (2021). Retrieved from [https://vanderbilt.policytech.com/Central Venous Access Devices: Insertion and Maintenance, and Discontinuation Universal Protocol - Identification of Correct Patient, Procedure, Site/Side \(Time Out\) CVAD: Insertion, Maintenance and Discontinuation – Children’s Hospital \(CL SOP\) CVAD: Insertion, Maintenance and Discontinuation for Vanderbilt University Adult Hospital and Clinics \(SOP\)](https://vanderbilt.policytech.com/Central-Venous-Access-Devices-Insertion-and-Maintenance-and-Discontinuation-Universal-Protocol-Identification-of-Correct-Patient-Procedure-Site/Side-Time-Out-CVAD-Insertion-Maintenance-and-Discontinuation-Childrens-Hospital-CL-SOP)
- Barsun, A., Sen, S., Palmieri, T. L., & Greenhalgh, D. G. (2014). Peripherally inserted central line catheter infections in burn patients. *J Burn Care Res*, 35(6), 514-517. <https://doi.org/10.1097/BCR.0000000000000045>
- Ciofi Silva, C. L., Rossi, L. A., Canini, S. R., Goncalves, N., & Furuya, R. K. (2014). Site of catheter insertion in burn patients and infection: a systematic review. *Burns*, 40(3), 365-373. <https://doi.org/10.1016/j.burns.2013.10.026>
- O'Grady, N. P., Alexander, M., Burns, L. A., Dellinger, E. P., Garland, J., Heard, S. O., Lipsett, P. A., Masur, H., Mermel, L. A., Pearson, M. L., Raad, I. I., Randolph, A. G., Rupp, M. E., & Saint, S. (2011). Guidelines for the Prevention of Intravascular Catheter-related Infections. *Clinical Infectious Diseases*, 52(9), e162-e193. <https://doi.org/10.1093/cid/cir257>
- O'Mara, M. S., Reed, N. L., Palmieri, T. L., & Greenhalgh, D. G. (2007). Central venous catheter infections in burn patients with scheduled catheter exchange and replacement. *J Surg Res*, 142(2), 341-350. <https://doi.org/10.1016/j.jss.2007.03.063>
- Practice Guidelines for Central Venous Access 2020: An Updated Report by the American Society of Anesthesiologists Task Force on Central Venous Access. (2020). *Anesthesiology*, 132(1), 8-43. <https://doi.org/10.1097/ALN.0000000000002864>
- Remington, L., Faraklas, I., Gauthier, K., Carper, C., Wiggins, J. B., Lewis, G. M., & Cochran, A. (2016). Assessment of a Central Line–Associated Bloodstream Infection Prevention Program in a Burn-Trauma Intensive Care Unit. *JAMA Surgery*, 151(5), 485. <https://doi.org/10.1001/jamasurg.2015.4445>
- Selby, L. M., Rupp, M. E., & Cawcutt, K. A. (2021). Prevention of Central-Line Associated Bloodstream Infections: 2021 Update. *Infect Dis Clin North Am*, 35(4), 841-856. <https://doi.org/10.1016/j.idc.2021.07.004>
- Ramos, G. E., Bolgiani, A. N., Patino, O., Prezzavento, G. E., Guastavino, P., Durlach, R., Fernandez Canigia, L. B., & Benaim, F. (2002). Catheter infection risk related to the distance between insertion site and burned area. *J Burn Care Rehabil*, 23(4), 266-271. <https://doi.org/10.1097/00004630-200207000-00007>