

VANDERBILT  UNIVERSITY
MEDICAL CENTER

Guideline: Early Excision and Grafting for Full Thickness
Burns >20% TBSA

Revised Date: July 2024

Review Date: July 2026

Content Experts

Robel Beyene, MD

Table of Contents

I.	Purpose	2
II.	Population	2
III.	Intervention/Treatment	2
IV.	Exceptions	3
V.	References	4

I. Purpose

Multiple studies have shown the benefit of Early Excision, variably defined as post-injury day 2 through 7, in patients with severe thermal injuries ($\geq 20\%$ Total Body Surface Area, TBSA). Outcomes affected by early excision and grafting include but are not limited to improved scarring and mobility, decreased rates of infection, shorter hospital length of stay, and improved survival rates. Burned skin results in massive systemic inflammatory response due the release of cytokines and growth factors. This inflammatory response is a large contributor to the fluid shifts that are seen in these patients and their fluid resuscitation requirements. This inflammatory response will continue until the burned skin is excised.

Additionally, devitalized skin is a rich medium for microbial overgrowth leading to infection. All burn wounds are colonized by bacteria and invasive wound sepsis has been greatly ameliorated by early excision and grafting, which removes the devitalized tissue before this colonization becomes infection. Early excision will now be tracked as a quality metric as time in days.

Early excision and grafting have been shown in multiple studies to modulate the systemic inflammatory response, attenuate muscle catabolism, and improve myocardial dysfunction caused by thermal injury. This protocol seeks to standardize our time frames for excision and grafting in our severely burned patients.

Ultra-early excision (within 2 days) has variably shown some benefits in weak data, such as fewer in-hospital complications. However, vasoactive use, blood loss, transfusion requirements, length of stay, readmission, and mortality are not clearly better than in the early groups. Some data suggests improvements in VTE/PE rates, CAUTI rates with ultra-early excision. Wounds excised within 3 days have some benefit regarding long term hypertrophic scar and contracture, but this must be achieved within the context of burn size and patient stability.

II. Population

All patients admitted to VUH and VCH with $\geq 20\%$ TBSA full thickness burns.

III. Intervention/Treatment

- a. Patients without contraindication (see exceptions below) will get their first excision of full thickness burns within the first 24-96 hours of their burn
- b. Pediatric patients should have their first excision within 24-48 hours and be fully excised within 24-96 hours
- c. Ideally, patients will reach the diuretic phase of acute burn resuscitation before initial excision, which generally occurs within the first 24 hours of resuscitation
- d. Patients will return to OR for excision and grafting (autograft vs. allograft vs. biologic matrix) every 24-72 hours until all full thickness burns are excised and grafted with autograft
 - i. If no further donor site is available, this process will stop when all full thickness burns are excised and grafted in skin substitute
 - ii. If the patient becomes unstable and is unable to return to the

operationroom within this timeframe, a discussion between the burn surgeon andburn intensivist will take place and a plan will be made and documentedin the medical record.

- e. Complete excision (>95%) of all full thickness burns should be complete within 7 days.
- f. If no further donor available and patient in skin substitute or dermal substitute:
- g. Patient will return to the OR every 7-10 days to exchange allograft/xenograft to prevent graft integration
- h. Any areas that are non-adherent either due to infection or non-viablewound bed will be excised within 24-72 hours of identification
- i. Patient will return to OR for autograft every 24-72 hours once donor sites areavailable again until grafting is complete
- j. Large TBSA burns may meet criteria for to be leveled to the operating room, based on physiologic changes secondary to the burn

IV. Exceptions/Contraindications

In patients that are unstable and cannot tolerate a trip to the operating room, a discussion between the burn surgeon and the burn intensivist will take place and a plan will be formulated and documented in the medical record.

V. References

1. Miroshnychenko A, Kim K, Rochweg B, Voineskos S. Comparison of early surgical intervention to delayed surgical intervention for treatment of thermal burns in adults: a systematic review and meta-analysis. *Burns Open*. 2021 Feb 25.
2. Herndon, D.N., *Total burn care*. 4th ed. 2012, Edinburgh ; New York: SaundersElsevier. xvii, 784 p.
3. Herndon, D.N., et al., *A comparison of conservative versus early excision. Therapies in severely burned patients*. *Ann Surg*, 1989. 209(5): p. 547-52; discussion 552-3.
4. Thompson, P., et al., *Effect of early excision on patients with major thermal injury*. *J Trauma*, 1987. 27(2): p. 205-7.
5. Herndon, D.N., et al., *Determinants of mortality in pediatric patients with greater than 70% full-thickness total body surface area thermal injury treated by early total excision and grafting*. *J Trauma*, 1987. 27(2): p. 208-12.
6. Horton, J.W., et al., *The effects of early excision and grafting on myocardial inflammation and function after burn injury*. *J Trauma*, 2006. 61(5): p. 1069- 77.
7. Gibran, N.S., et al., *Summary of the 2012 ABA Burn Quality Consensus conference*. *J Burn Care Res*, 2013. 34(4): p. 361-85.
8. Barret, J.P. and D.N. Herndon, *Modulation of inflammatory and catabolic responses in severely burned children by early burn wound excision in the first 24 hours*. *Arch Surg*, 2003. 138(2): p. 127-32.
9. Hayashi K, Sasabuchi Y, Matsui H, et al. Does early excision or skin grafting of severe burns improve prognosis? A retrospective cohort study. *Burns*. 2023 May 1;49(3):554-61.
10. De La Tejera G, Corona K, Efejuku T, et al. Early wound excision within three days decreases risks of wound infection and death in burned patients. *Burns*. 2023 Dec 1;49(8):1816-22.
11. Ramsey WA, O'Neil Jr CF, Corona AM, et al. Burn excision within 48 hours portends better outcomes than standard management: A nationwide analysis. *JTACS*. 2023 Jul 1;95(1):111-5.