

The Vanderbilt Child Health poll surveyed 1,026 Tennessee parents with children under 18 years old from October 19 —November 28, 2021. All reported percentages have survey weights applied, so their interpretation can be broadened to all parents in Tennessee. For example, “13.2% of Tennessee parents are concerned their child might have Depression that has not yet been diagnosed by a medical professional.”

All summaries are listed as weighted proportions of Tennessee Parents ± Standard of Error.

### CHILDREN’S BEHAVIORAL HEALTH

*If you have a child aged 6-17, has your child been **diagnosed** by a medical professional with any of the following conditions?*

*If you have a child aged 6-17, are you concerned your child might have one of the following conditions, but **has not yet been diagnosed**?*

\*It was possible to choose more than one answer

	2021	2020	2019	2021	2020	2019
<b>ADHD/ADD</b>	15.0% ± 1.7	11.1% ± 1.4	15.7% ± 1.8	7.6% ± 1.2	6.7% ± 1.3	6.7% ± 1.3
<b>Anxiety</b>	15.5% ± 1.8	9.8% ± 1.2	10.9% ± 1.4	17.1% ± 1.9	21.7% ± 2.2	13.8% ± 1.5
<b>Autism</b>	3.6% ± 0.9	4.9% ± 1.0	NA	1.8% ± 0.6	1.9% ± 0.6	NA
<b>Bipolar disorder</b>	1.2% ± 0.3	3.0% ± 0.8	2.0% ± 0.5	5.3% ± 1.2	5.5% ± 1.1	4.7% ± 1.0
<b>Depression</b>	7.9% ± 1.1	7.2% ± 1.2	7.5% ± 1.2	13.2% ± 1.7	10.6% ± 1.5	11.8% ± 1.7
<b>Eating disorder</b>	3.1% ± 0.8	3.3% ± 0.7	1.1% ± 0.5	3.0% ± 0.7	3.9% ± 1.1	3.1% ± 1.0
<b>ODD</b>	1.0% ± 0.4	1.6% ± 0.5	2.0% ± 0.6	2.6% ± 0.6	2.5% ± 0.8	2.5% ± 0.8
<b>PTSD</b>	1.6% ± 0.5	2.8% ± 0.6	2.3% ± 0.5	3.6% ± 1.1	5.3% ± 1.2	2.4% ± 0.7
<b>Self-harm</b>	2.3% ± 0.7	3.5% ± 0.8	3.0% ± 0.9	1.2% ± 0.4	2.2% ± 0.6	2.4% ± 0.6
<b>Stress</b>	5.6% ± 0.9	6.5% ± 1.1	4.2% ± 0.8	16.0% ± 1.9	11.6% ± 1.6	10.4% ± 1.4
<b>Substance use disorder</b>	1.1% ± 0.5	1.4% ± 0.6	1.1% ± 0.6	2.6% ± 0.6	2.4% ± 0.6	1.7% ± 0.6
<b>Suicidal thoughts/intentions</b>	2.6% ± 0.7	3.4% ± 0.8	2.4% ± 0.7	4.0% ± 1.0	2.9% ± 0.7	3.8% ± 0.9
<b>Other</b>	0.7% ± 0.3	2.8% ± 0.8	2.8% ± 0.7	0.1% ± 0.1	0.5% ± 0.2	0.1% ± 0.1
<b>None of these</b>	67.0% ± 2.3	68.3% ± 2.3	70.6% ± 2.2	60.4% ± 2.4	58.4% ± 2.6	66.1% ± 2.3

**STRATIFIED BY RACE/ETHNICITY GROUPS AND TENNESSEE REGIONS**

<b><i>Diagnosed Condition</i></b>	<b>N</b>	<b>Anxiety</b>	<b>Stress</b>	<b>Depression</b>	<b>ADHD/ADD</b>
<b>Race/Ethnicity</b>					
White, Non-Hispanic	625	16.0% ± 1.9	6.1% ± 1.0	9.1% ± 1.4	14.5% ± 1.8
Black, Non-Hispanic	93	9.8% ± 5.0	1.9% ± 1.0	2.1% ± 1.5	14.5% ± 5.4
Other/2+ races, Non-Hispanic/Hispanic	85	20.6% ± 7.5	7.9% ± 3.8	8.5% ± 4.2	20.2% ± 6.6
<b>TN Region</b>					
West	168	13.1% ± 4.1	4.4% ± 1.5	5.3% ± 1.9	11.7% ± 4.0
Middle	295	15.3% ± 2.8	5.2% ± 1.5	9.5% ± 2.1	15.6% ± 2.9
East	339	16.9% ± 2.8	6.5% ± 1.5	7.8% ± 1.6	16.3% ± 2.6

  

<b><i>Undiagnosed Condition</i></b>	<b>N</b>	<b>Anxiety</b>	<b>Stress</b>	<b>Depression</b>	<b>ADHD/ADD</b>
<b>Race/Ethnicity</b>					
White, Non-Hispanic	627	18.4% ± 2.3	15.4% ± 2.1	11.9% ± 1.7	8.8% ± 1.5
Black, Non-Hispanic	93	10.0% ± 3.1	16.0% ± 5.8	16.3% ± 5.1	3.5% ± 1.7
Other/2+ races, Non-Hispanic/Hispanic	84	19.9% ± 6.2	20.3% ± 6.5	17.5% ± 6.3	5.3% ± 3.2
<b>TN Region</b>					
West	169	15.7% ± 3.5	24.3% ± 5.5	13.0% ± 3.4	6.6% ± 2.1
Middle	295	21.7% ± 3.6	15.5% ± 2.8	14.7% ± 3.1	8.9% ± 2.3
East	339	13.0% ± 2.2	11.7% ± 2.5	11.7% ± 2.1	6.9% ± 1.6

Note: N's are unweighted; row-wise percentages presented (e.g. "16.3% of Black, Non-Hispanic parents were worried about undiagnosed Depression in their children, compared to 11.9% of White, Non-Hispanic parents")

\* No statistically significant differences across categories by Rao-Scott corrected Chi-Square testing