

CATHETER-ASSOCIATED URINARY TRACT INFECTION (CAUTI) PREVENTION BUNDLE

INSERTION OF CATHETER

- ◆ Use Aseptic Technique
 - ◆ Hand hygiene
 - ◆ Prior to establishing sterile field
 - ◆ Prior to donning gloves and completing insertion
 - ◆ After discarding supplies and removing gloves
 - ◆ Catheter Insertion Kit
 - ◆ Sterile gloves, PPE if necessary
 - ◆ Pre-cleanse kit
 - ◆ Povidone/iodine prep
 - ◆ New catheter and sterile gloves with each unsuccessful attempt
- ◆ Avoid Unnecessary Catheterizations
 - ◆ Written order prior to placement or within 24 hours in emergent situations

MAINTENANCE OF CATHETER

- ◆ Maintain Closed Drainage System
 - ◆ Minimal access with aseptic technique
 - ◆ Hand hygiene and clean gloves
 - ◆ CHG scrub to disinfect access port
 - ◆ Drainage port clean and clamped and not touching receiving container
- ◆ Maintain Hygiene
 - ◆ Perineal/meatal care every 12 hours and PRN
- ◆ Keep Bag Below Level of Bladder
 - ◆ Drainage system above floor level and below bladder level
- ◆ Maintain Unobstructed Urine Flow
 - ◆ Empty collection bag every 12 hours, before transport and PRN (be sure to measure and document output every 2 hours)
- ◆ Keep Catheter Anchored at All Times
- ◆ Remove Catheter When No Longer Needed
 - ◆ Daily discussion of need and functionality

