TASSO-SST

Button

VIDEO

PREPARE

FIRST TIMER USERS Point your phone's camera here to watch:



or go to www.tassoinc.com/ tasso-sst-video




1. Get a timer. You'll need it in step 9.



2. Rub arm quickly and firmly just below the shoulder.

Rub until it's very warm to help your blood flow.



3. Clean arm with alcohol pad.



4. Open device pouch by pulling apart white and clear layers.

COLLEC



5. Remove clear plastic clover over the red button.



6. Peel paper tab behind the red button.

Keep tube pointing down.



7. Stick device to shoulder.

Do not remove once it's on.



8. Press button quickly and firmly until it can't go any farther. Wait two seconds then let go.



9. Start a five minute timer.

Keep arm at your side. You won't see blood right away.



10. After five minutes, peel off device.

No need to fill the tube completely.

RETURN



11. Remove tube by firmly twisting a quarter turn and pulling down.

This may take a bit of finger strength.





12. Twist cap fully onto the tube.



13. Place tube in the clear return bag and seal.

Leave the absorbent pad in the bag.



14. Write collection date and time on the inside flap of the box.



15. Place bag in box and seal with the strip.



16. Return box within24 hours of packing.

Throw out used Button in standard household trash.

For in vitro diagnostic use IND Lot number TOJ Catalog number REF Sterilized by irradiation STERILE R **SAMBOLS** Prescription use only **BX ONLY** Store at 15-30°C Instructions for Use (\$ Return Pouch with Absorbent Pad Vino seu signis Bandage Do not use if package is damaged (8) Alcohol Wipe $\overline{\mathbb{V}}$ This device contains sharps Cap for Collection Tube Consult instructions for use Tasso Blood Collection Device Use by date KIT CONTENTS

> shipped back to a laboratory for measurement of COVID-19 antibodies in blood. self-collection of capillary blood from the upper arm of adults. The sample is TASSO-SST is a single use blood collection device that is intended for the


Use while seated as fainting may occur with any blood sampling procedure.

Intended for self-use or use by a healthcare provider.

Use before the expiration date affixed to the box.

Minor bruising may occur around the sample collection site.

PN0101-01, A2





INSTRUCTIONS FOR USE AGES 4 AND OLDER



TASSO ONDEMAND **TASSO-SST**

Contact us for help: 1-408-708-8451

INTENDED USE

For external use only.

One-time use only.

Prescription use only.

Keep out of reach of children.