

EMERGENCY GENERAL SURGERY

Practice Management Guideline: Percutaneous Endoscopic Gastrostomy (PEG) Tube

- I. Background:** Percutaneous endoscopic gastrostomy (PEG) is a commonly performed procedure that provides durable enteral access for feeding and medications. This document outlines patient selection and perioperative considerations for patients undergoing PEG tube placement.
- II. Guideline:**
- A. Patient Selection/Preoperative Care**
- a. Contraindications to PEG Placement
 - i. Anatomic inability to place PEG (e.g. Roux-en-Y anatomy, mesh overlying PEG site, esophageal stricture)
 - ii. Uncontrolled agitation (including any patient requiring arm restraints or sitter for agitation in prior 48 hours)
 - iii. Ascites
 - iv. Severe Dementia
 - v. Anorexia Nervosa
 - b. Special Situations for PEG Placement
 - i. Esophageal cancer: discuss with Thoracic Surgery to ensure no plans for esophagectomy with gastric conduit
 - ii. Failure to thrive: will perform if there is a medically reversible cause
 - iii. Palliative Decompression: In the setting of unresectable malignancy. Should be pursued only with input of palliative care team and/or surgical oncology team
 - iv. Peritoneal Dialysis (PD): Patients must be on hemodialysis for 6 weeks prior to PEG and wait at least 4 weeks to go back to PD
 - v. Ventriculoperitoneal (VP) shunt: VP shunt and PEG placement should be separated by 7 days to reduce risk of infectious complications
 - vi. Amyotrophic Lateral Sclerosis (ALS) *without* tracheostomy: Interventional Radiology (IR)-placed gastrostomy preferred to PEG given the anesthesia risk in this population
 - vii. Active chemotherapy
 1. Risk/benefit discussion of immediate vs. delayed PEG
 2. Contraindicated during severe leukopenia (WBC <2.5 or ANC 0.5) or thrombocytopenia (<50)
 - viii. Immunosuppressants/Corticosteroids
 1. Risk/benefit discussion of immediate vs. Delayed PEG with medical oncology team
 2. Contraindicated during induction dose
 3. If plans to stop steroids, delay until no longer on steroids
 4. If no plans to stop steroids, delay until on maintenance/long term dose
 - c. Anticoagulation Hold Times
 - i. Heparin drip: Hold 4 hours prior to and 4 hours after procedure
 - ii. Prophylactic dose enoxaparin or heparin: do not hold
 - iii. Therapeutic enoxaparin: hold morning dose, restart evening dose if no bleeding
 - iv. Direct Oral Anticoagulants (DOAC): Hold 24 hours prior to PEG, restart evening dose if no bleeding
 - v. Aspirin: Do not hold

- vi. Antiplatelet agents, e.g. Clopidogrel/Ticagrelor/Effient (including in setting of dual antiplatelet therapy): Do not hold
- d. Preoperative Tube Feeds and Diet
 - i. Bedside procedures (in ICU)
 - 1. Protected airway (cuffed tracheostomy or endotracheal tube): hold feeds minimum of one hour. *This includes patients undergoing percutaneous tracheostomy at the same time.*
 - 2. Unprotected airway: Hold tube feeds/solid foods minimum of 6 hours, clear liquids 2 hours
 - ii. Operating room procedures:
 - 1. Hold tube feeds and diet at midnight the night prior. For patients at high nutritional risk hold solid foods/tube feeds minimum 6 hours, clear liquids minimum 2 hours.

B. Procedural Care

- a. Follow bedside surgery protocol for sterile precautions, informed consent, sedation, and preoperative timeout
- b. Equipment: Refer to **Appendix A**
- c. Antibiotics: Weight-based Cefazolin given within 1 hour of incision
- d. Indications for T-fasteners:
 - i. Corticosteroids
 - ii. Chemotherapy
 - iii. Immunosuppressants
 - iv. Attending discretion

C. Postoperative Care

- a. Postoperative PEG Care
 - i. PEG to gravity drainage (foley bag) x 4 hours
 - ii. Tube feeds and medications started after 4 hours
 - iii. Patients are followed with a post-op check and note and a POD1 visit and note with depth documented
 - iv. Those who are higher risk or have additional identified issues should continue to be followed
 - v. T fasteners removed at 2-3 weeks unless causing skin necrosis
 - vi. Post-discharge PEG appointment made for 4 weeks following placement
 - vii. Dislodged PEG should have urgent EGS evaluation
- b. Bedside Nursing Care: **Appendix B**
- c. Dislodged PEG/Gastrostomy Tube
 - i. Dislodged tube within 30 days should have emergent EGS consult/evaluation
 - ii. Peritonitis, free fluid or air, hemodynamic changes, abdominal wall infection necessitate consideration of exploration
 - iii. If no peritonitis, attempt replacement with balloon gastrostomy tube, confirm replacement with either X-ray tube study or IV and per-tube contrasted CT scan
 - iv. Consider repeat PEG in patients without evidence of peritonitis or abdominal wall infection
- d. PEG Removal
 - i. PEG should not be removed for a *minimum of 6 weeks* post-placement

- ii. PEG Removal Criteria:
 - 1. Initial pathology improved (i.e. cleared by SLP for PO diet)
 - 2. All food should be taken PO only for 2 weeks
 - 3. Patient's weight stable on PO diet
 - 4. Tolerating multiple consistencies of foods, good appetite, not overly reliant on supplements
 - 5. No pending surgery or radiation planned
- iii. High dose PPI may be used to facilitate gastrocutaneous fistula closure

III. References

- Abraham, N.S., et al., *American College of Gastroenterology-Canadian Association of Gastroenterology Clinical Practice Guideline: Management of Anticoagulants and Antiplatelets During Acute Gastrointestinal Bleeding and the Periendoscopic Period*. J Can Assoc Gastroenterol, 2022. **5**(2): p. 100-101.
- Bischoff, S.C., et al., *ESPEN practical guideline: Home enteral nutrition*. Clin Nutr, 2022. **41**(2): p. 468-488.
- Boullata, J.I., et al., *ASPEN Safe Practices for Enteral Nutrition Therapy [Formula: see text]*. JPEN J Parenter Enteral Nutr, 2017. **41**(1): p. 15-103.
- Committee, A.S.o.P., et al., *The management of antithrombotic agents for patients undergoing GI endoscopy*. Gastrointest Endosc, 2016. **83**(1): p. 3-16.
- Committee, A.S.o.P., et al., *Antibiotic prophylaxis for GI endoscopy*. Gastrointest Endosc, 2015. **81**(1): p. 81-9.
- Davies, N., et al., *Enteral tube feeding for people with severe dementia*. Cochrane Database Syst Rev, 2021. **8**(8): p. CD013503.
- de Sousa Magalhaes, R., et al., *Percutaneous endoscopic gastrostomy: dealing with the issue of dislodgement*. Scand J Gastroenterol, 2020. **55**(4): p. 485-491.
- Elfert, K et al. *Impact of antithrombotic medications on postprocedural outcomes of percutaneous endoscopic gastrostomy: a US Collaborative Network study*. European Journal of Gastroenterology & Hepatology, 2025. **37** (9), 1021-1025.
- Gangwani, M.K., et al., *Dual Antiplatelet Therapy Does Not Increase Bleeding Risk in Percutaneous Gastrostomy Tube Placement: Network Meta-Analysis*. Dig Dis Sci, 2023. **68**(5): p. 1966-1974.
- Keung, E.Z., et al., *In-hospital and long-term outcomes after percutaneous endoscopic gastrostomy in patients with malignancy*. J Am Coll Surg, 2012. **215**(6): p. 777-86.
- Lipp, A. and G. Lusardi, *Systemic antimicrobial prophylaxis for percutaneous endoscopic gastrostomy*. Cochrane Database Syst Rev, 2013. **2013**(11): p. CD005571.
- Lucendo, A.J., et al., *Risk of bleeding in patients undergoing percutaneous endoscopic gastrostomy (PEG) tube insertion under antiplatelet therapy: a systematic review with a meta-analysis*. Rev Esp Enferm Dig, 2015. **107**(3): p. 128-36.
- Meenaghan, N., K. Lumpkins, and J. Scott Roth, *Percutaneous endoscopic gastrostomy tube placement is safe in patients undergoing corticosteroid therapy*. J Gastrointest Surg, 2009. **13**(2): p. 236-8.
- Oterdoom, L.H., et al., *Systematic review of ventricular peritoneal shunt and percutaneous endoscopic gastrostomy: a safe combination*. J Neurosurg, 2017. **127**(4): p. 899-904.
- Rosenberger, L.H., et al., *Late accidental dislodgement of a percutaneous endoscopic gastrostomy tube: an underestimated burden on patients and the health care system*. Surg Endosc, 2011. **25**(10): p. 3307-11.

IV. Authors

Elizabeth D. Krebs, MD
Michael C. Smith, MD
Nina E. Collins, APRN

May 1, 2024, Updated May 15, 2026

Appendix A: Procedural Equipment Needed for PEG

1. Trauma procedure cart
2. Sterile towels
3. Bite Block
4. Sterile Gown and Gloves
5. Eye Protection
6. Scope
 - Pediatric Colonoscope for PEG-J
 - Pediatric XP scope for restricted esophagus
7. 1-liter bottle normal saline
8. Endoscopy connector set
9. Suction Tubing and Canister
10. Chlorhexidine skin prep
11. PEG or PEG-J kit (If Planning PEG/J T-fasteners are required).

Appendix B: Bedside Gastrostomy Care

- 1) General care of gastrostomy tube
 - i) Record length of the tube where bumper base is located every shift.
 - ii) Cleanse site every shift with NS moist gauze to remove crust or drainage.
 - (a) If breakdown or macerated skin at the gastrostomy tube entrance site, apply Triple Care EPC and dry gauze daily. Change gauze PRN.
 - (b) If no breakdown and skin is intact, apply Triple Care Ointment or Ilex Cream and dry gauze daily.
 - iii) Change gauze PRN.
 - iv) Flush tube with 30 milliliters of water every 8 hours and PRN after giving medication or feedings per the tube.
 - v) Apply abdominal binder or appropriate tape to keep gastrostomy tube from hanging out of patient clothing or to prevent unintended dislodgement of tube.
- 2) PEG tube feedings, supplements, normal saline, and water are to be given as directed by the attending physician or per nutrition services as directed in the orders.
- 3) Any dislodged tube or tube with differing tube lengths compared to prior should have tube feeds stopped and EGS team notified