



VANDERBILT HEALTH
DIVISION OF ACUTE CARE SURGERY

Perioperative Antibiotic Prophylaxis in Trauma (Non-Orthopedic)



For emergent chest tube placement or operative cases, the order should be entered as “stat”. However, for non-emergent chest tube placement or operative cases, cefazolin should be ordered to be given 30 minutes prior to incision (within 120 minutes for levofloxacin given prolonged infusion time)



**Patients receiving blood transfusions of 10 units or greater may require repeat antimicrobial dosing intraoperatively. Consider re-dosing of ceftriaxone or levofloxacin x1 post-operatively if >2 coolers massive transfusion received and antibiotics were not re-dosed during OR.

- Prophylactic antifungal coverage is not required for upper GI injuries if repaired within 24-48 hours.
- Extended therapy is not needed with gross contamination or in cases of penetrating injuries to the thoracolumbar spine with associated intraperitoneal injuries.¹

RATIONALE

It is important to first determine whether to pursue prophylactic antibiotic coverage or treatment of active infection. If true infection is suspected, patients should be managed in accordance to the Infectious Workup and Antimicrobial Stewardship Protocol. If prophylactic coverage is desired, the above protocol should be followed.

There is some controversy surrounding the use of prophylactic antibiotics in patients receiving chest tubes, however, adequate data exist to support their use. A single dose of narrow, gram positive coverage is warranted.³⁻⁵

Antibiotic therapy should be continued for **no longer than 24 hours**, as prophylaxis beyond 24 hours is not associated with a risk of surgical infections, but rather an increase in nosocomial infectious complications and resistant infections.^{2, 6-7}

The post-operative infection rate following surgery for abdominal trauma with hollow viscus injury is between 10 and 15%.⁸ Demonstrated risk factors for infection include hollow viscus injury, presence and degree of intestinal contamination, patient age, transfusion requirement, and shock. The microbiology and bacterial load differ throughout the gastrointestinal tract; the concentration of bacteria increases in proximal to distal direction.

Broad spectrum, prophylactic antibiotic coverage should be directed against gram positive, enteric gram-negative and anaerobic bacteria.⁹ Common pathogens include:

- Streptococcus
- Enterococcus
- Enterobacteriaceae
- Bacteroides
- Peptostreptococcus
- Clostridium

Many regimens have been proven to be efficacious in this setting. Second-generation cephalosporins and extended spectrum penicillins are as effective as regimens which include aminoglycosides and in addition, offer the convenience of single agent therapy. Aminoglycosides are no longer recommended due to suboptimal activity in this patient population. Agents whose spectra include coverage of enterococcus species may be more efficacious than those that do not.¹⁰

REFERENCES

1. Pasupuleti LV, Sifri ZC, Mohr AM. Is Extended Antibiotic Prophylaxis Necessary after Penetrating Trauma to the Thoracolumbar Spine with Concomitant Intraperitoneal Injuries? *Surg Infect (Larchmt)*. 2014;15(1):8-13.
2. Goldberg SR, Anand RJ, Como JJ et al. Prophylactic antibiotic use in penetrating abdominal trauma: An Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg*. 2012;75(5):321-325.
3. Moore FO, Duane TM, Hu CK et al. Presumptive antibiotic use in tube thoracostomy for traumatic hemopneumothorax: An Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg*. 2012;73(5):S341-S344.
4. Bosman A, de Jong MB, Debeij J et al. Systematic review and meta-analysis of antibiotic prophylaxis to prevent infections from chest drains in blunt and penetrating thoracic injuries. *Br J Surg* 2012;99:506-513.
5. Heydari MB et al. Use of prophylactic antibiotics following tube thoracostomy for blunt chest trauma in the prevention of empyema and pneumonia. *J Inj Violence Res* 2014;6(2):91-92.
6. Fabian TC, Croce MA, Payne LW et al. Duration of antibiotic therapy for penetrating abdominal trauma: a prospective trial. *Surgery* 1992; 112:788-794.
7. Velmahos GC, Toutouzas KG, Sarkisyan G et al. Severe trauma is not an excuse for prolonged antibiotic prophylaxis. *Arch Surg* 2002; 137:537-541.
8. Fabian TC. Infection in penetrating abdominal trauma: Risk factors and preventive antibiotics. *Am Surg*. 2002;68(1):29-35.
9. Fabian TC. Prevention of infections following penetrating abdominal trauma. *Am J Surg* 1993; 165:14S-19S.
10. Weigelt JA, Easley SM, Thal ER et al. Abdominal surgical wound infection is lowered with improved perioperative enterococcus and bacteroides therapy. *J Trauma* 1993; 34:579-584.
11. Huston JM, Kreiner L, Ho VP, et al. Role of Empiric Anti-Fungal Therapy in the Treatment of Perforated Peptic Ulcer Disease: Review of the Evidence and Future Directions. *Surg Infect (Larchmt)*. 2019;20(8):593-600.
12. Barmparas G, Alhaj Saleh A, Huang R, et al. Empiric antifungals do not decrease the risk for organ space infection in patients with perforated peptic ulcer. *Trauma Surg Acute Care Open*. 2021;6(1):e000662.
13. Bratzler DW, Dellinger EP, Olsen KM, et al; American Society of Health-System Pharmacists; Infectious Disease Society of America; Surgical Infection Society; Society for Healthcare Epidemiology of America. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Am J Health Syst Pharm*. 2013 Feb 1;70(3):195-283.
14. Farrell MS, Agapian JV, Appelbaum RD, et al. Surgical and procedural antibiotic prophylaxis in the surgical ICU: an American Association for the Surgery of Trauma Critical Care Committee clinical consensus document. *Trauma Surg Acute Care Open*. 2024 Jun 3;9(1):e001305.

Revised July 2026

Bradley Dennis, MD

Jill Streams, MD

Jennifer Beavers, PharmD

Chelsea Tasaka, PharmD