


Diabetes Insipidus (DI) Following Traumatic Brain Injury



During therapy: Check electrolytes at least q4 hours in the acute phase, follow urine specific gravity, strict I&Os, and be aware that sodium should **NOT** rise or fall > 1mEq/h

References

- 1 Anders, Brian T. Intensive Care in Neurosurgery. New York: American Association of Neurological Surgeons, 2003, pp 66-69.
- 2 Oiso Y, Robertson GL, Norgaard JP, Juul, KV. Treatment of neurohypophyseal diabetes insipidus. J Clin Endocrinol Metab. 2013;98:3958-3967.
- 3 Qureshi S, Galiveeti S, Bichet DG, Roth J. Diabetes Insipidus: Celebrating a century of vasopressin therapy. Endocrinology 2014;155:4605-4621.
- 4 Shucart WA, Jackson I. Management of diabetes insipidus in neurosurgical patients. J Neurosurg. 1976;44:65-71.
- 5 Marino P. The ICU Book: 4th Edition. New York: Wolters Kluwer/Lippincott Williams and Wilkins, 2014, pp 661-663.

Updated/Revisions:

May 2020, June 2022, May 2024

Authors:

Brad Dennis, MD

Laurie B Ford, ACNP-BC

Joshua P Smith, DO

Leanne Atchison, PharmD

Jennifer Beavers, PharmD