


Hemothorax Guidelines

1. Place chest tube if HTX present per chest tube protocol (1)
 - a. Pigtail or large bore chest tube (14-32F) (2-4)
 - b. One dose of antibiotics starting just prior to placement (5, 6)
 - i. 1st generation cephalosporin
2. Order post-procedure CXR
3. If radiographic evidence of retained HTX
 - a. **If within 24 hours**, consider placing 2nd chest tube
 - b. Order a **contrasted chest CT on day 3** if HTX is still present on CXR (7)
 - i. If $\geq 300\text{mL}$ HTX (formula $v=d^2 \times l$) ¹ present on CT (8, 9)
 1. VATS on or before day 7 (1, 7-13)
 2. Consider intrapleural -tPA (11, 14-18) if poor operative candidate or other indications (t-PA is contraindicated in any patient with intrathoracic arterial bleeding/injury including pulmonary laceration and/or intercostal bleeding). Administration of TPA is not intended to prevent VATS in operative candidates.²
 - a. 24mg t-PA in 48mL NS (t-PA concentration 0.5mg/mL) injected in chest tube daily \times 3 days (16, 19-21) using sterile technique
 - b. Clamp chest tube \times 1 hour (20)
 - c. Roll patient to ensure distribution throughout chest
 - d. Unclamp tube and allow drainage
 - ii. If $< 300\text{ mL}$ HTX present, no additional intervention needed (1, 8)
 4. If no radiographic evidence of retained HTX
 - a. Place tube to water seal if no evidence of PTX on morning CXR and no air leak in tube
 - b. Remove chest tube when output $< 200\text{ mL}/24\text{hr}$ (22)

¹ **d** = greatest depth of hemothorax on a single CT image, **l** = greatest length of the hemothorax.

² To find the Trauma Retained Hemothorax order set in eStar you must type “hemothorax” in the order section, order set will then populate for you. It will not come up if you type t-PA or alteplase.



DIVISION OF TRAUMA AND SURGICAL CRITICAL CARE

1. Mowery NT, Gunter OL, Collier BR, et al. Practice management guidelines for management of hemothorax and occult pneumothorax. *J Trauma*. 2011 Feb;70(2):510-8.
2. Inaba K, Lustenberger T, Recinos G, et al. Does size matter? A prospective analysis of 28-32 versus 36-40 French chest tube size in trauma. *J Trauma Acute Care Surg*. 2012 Feb;72(2):422-7.
3. Bauman ZM, Kulvatunyou N, Joseph B, et al. Randomized Clinical Trial of 14-French (14F) Pigtail Catheters versus 28-32F Chest Tubes in the Management of Patients with Traumatic Hemothorax and Hemopneumothorax. *World J Surg*. 2021 Mar;45(3):880-6.
4. Kulvatunyou N, Bauman ZM, Edine SBZ, et al. The Small 14-French (Fr) Percutaneous Catheter vs. Large (28-32Fr) Open Chest Tube for Traumatic Hemothorax (P-CAT): A Multi-center Randomized Clinical Trial. *J Trauma Acute Care Surg*. 2021 Mar 16.
5. Luchette FA, Barrie PS, Oswanski MF, et al. Practice Management Guidelines for Prophylactic Antibiotic Use in Tube Thoracostomy for Traumatic Hemopneumothorax: the EAST Practice Management Guidelines Work Group. *Eastern Association for Trauma. J Trauma*. 2000 Apr;48(4):753-7.
6. Maxwell RA, Campbell DJ, Fabian TC, et al. Use of presumptive antibiotics following tube thoracostomy for traumatic hemopneumothorax in the prevention of empyema and pneumonia--a multi-center trial. *J Trauma*. 2004 Oct;57(4):742-8; discussion 8-9.
7. Velmahos GC, Demetriades D, Chan L, et al. Predicting the need for thoracoscopic evacuation of residual traumatic hemothorax: chest radiograph is insufficient. *J Trauma*. 1999 Jan;46(1):65-70.
8. DuBose J, Inaba K, Demetriades D, et al. Management of post-traumatic retained hemothorax: a prospective, observational, multicenter AAST study. *J Trauma Acute Care Surg*. 2012 Jan;72(1):11-22; discussion -4; quiz 316.
9. Mergo PJ, Helmberger T, Didovic J, Cerniglio J, Ros PR, Staab EV. New formula for quantification of pleural effusions from computed tomography. *J Thorac Imaging*. 1999 Apr;14(2):122-5.
10. Oguzkaya F, Akcali Y, Bilgin M. Videothoracoscopy versus intrapleural streptokinase for management of post traumatic retained haemothorax: a retrospective study of 65 cases. *Injury*. 2005 Apr;36(4):526-9.
11. Boersma WG, Stigt JA, Smit HJ. Treatment of haemothorax. *Respir Med*. 2010 Nov;104(11):1583-7.
12. Heniford BT, Carrillo EH, Spain DA, Sosa JL, Fulton RL, Richardson JD. The role of thoracoscopy in the management of retained thoracic collections after trauma. *Ann Thorac Surg*. 1997 Apr;63(4):940-3.
13. Meyer DM, Jessen ME, Wait MA, Estrera AS. Early evacuation of traumatic retained hemothoraces using thoracoscopy: a prospective, randomized trial. *Ann Thorac Surg*. 1997 Nov;64(5):1396-400; discussion 400-1.
14. Agarwal R, Aggarwal AN, Gupta D. Intrapleural fibrinolysis in clotted haemothorax. *Singapore Med J*. 2006 Nov;47(11):984-6.
15. Inci I, Ozcelik C, Ulku R, Tuna A, Eren N. Intrapleural fibrinolytic treatment of traumatic clotted hemothorax. *Chest*. 1998 Jul;114(1):160-5.
16. Ben-Or S, Feins RH, Veeramachaneni NK, Haithcock BE. Effectiveness and risks associated with intrapleural alteplase by means of tube thoracostomy. *Ann Thorac Surg*. 2011 Mar;91(3):860-3; discussion 3-4.
17. Stiles PJ, Drake RM, Helmer SD, Bjordahl PM, Haan JM. Evaluation of chest tube administration of tissue plasminogen activator to treat retained hemothorax. *Am J Surg*. 2014 Jun;207(6):960-3.
18. Constantine B, Hobbs B, Cheatham M. Tissue Plasminogen Activator in Traumatic Hemothorax. Available at: <http://surgicalcriticalcare.net/Guidelines/Tissue%20plasminogen%20activator%20in%20traumatic%20hemothorax%202018.pdf>. Accessed June 2, 2021.
19. Skeete DA, Rutherford EJ, Schlidt SA, Abrams JE, Parker LA, Rich PB. Intrapleural tissue plasminogen activator for complicated pleural effusions. *J Trauma*. 2004 Dec;57(6):1178-83.

DIVISION OF TRAUMA AND SURGICAL CRITICAL CARE

20. Thommi G, Nair CK, Aronow WS, Shehan C, Meyers P, McLeay M. Efficacy and safety of intrapleural instillation of alteplase in the management of complicated pleural effusion or empyema. *Am J Ther.* 2007 Jul-Aug;14(4):341-5.
21. Hamblin SE, Furmanek DL. Intrapleural tissue plasminogen activator for the treatment of parapneumonic effusion. *Pharmacotherapy.* 2010 Aug;30(8):855-62.
22. Younes RN, Gross JL, Aguiar S, Haddad FJ, Deheinzelin D. When to remove a chest tube? A randomized study with subsequent prospective consecutive validation. *J Am Coll Surg.* 2002 Nov;195(5):658-62.

Revisions:

June 2018

June 2021

Bradley M. Dennis, MD

Susan Hamblin PharmD

Bethany Evans, ACNP-BC