

**Surgical Intensive Unit
Guidelines for the Mobilization of the Critically Ill Patient**

I. Purpose


To utilize a multi-disciplinary approach to the mobilization of critically ill patients in the Surgical Intensive Care Unit

II. Process

Upon the start of each shift, the nurse will refer to the Safety Screen below to evaluate the patient’s appropriateness for mobilization.

- Neuro: Brain death, ICP >15, active seizure, neuromuscular blockade, spinal cord injury
- Cardiovascular: Infusion of two or more vasopressors, MAP <65, SBP <90 or >170, active myocardial infarction or unstable arrhythmia
- Respiratory: Unstable airway, Ventilator FiO2 >70%, PEEP > 15, Bi-vent or VDR settings
- Gastrointestinal: Open abdomen
- Miscellaneous: Comfort Care, presence of femoral sheath, order for bedrest

The nurse will communicate the pass or fail of the Safety Screen during multi-disciplinary rounds. The Critical Care team during rounds will formulate a mobility goal for the patient based on the pyramid below.



Starting at Level One, the bedside team will progress the patient in a step-like manner up the pyramid to the goal established during rounds. If the patient develops a limitation to mobilization, including, but not limited to, hemodynamic instability, nausea, or pain control, the bedside team will assist the patient to the prior level of mobility, notify the critical care team of the limitation, and re-attempt mobility after 30 minutes of rest.

III. Lead Author

Carolyn Ragland, DNP, AGACNP-BC
Critical Care Nurse Practitioner

Michael C. Smith, MD
SICU PI Physician Lead

IV. Endorsement

Shannon Eastham, MD
SICU Co-Medical Director

Revised June 2021