
ORTHOPEDIC PROPHYLACTIC ANTIBIOTIC PROTOCOLS

Rationale:

The rate of infections in open fractures is significantly related with the degree of soft tissue injury, contamination, blood supply to the area, and to the management of the wound¹⁻³. Most commonly open fractures are classified by the Gustilo classification system:

Type I	wound < 1 cm in length, clean with minimal soft tissue loss or injury
Type II	wound > 1 cm but <10 cm without extensive soft tissue injury, tissue loss
Type III	wound >10 cm with extensive soft tissue injury
	A: Adequate soft tissue coverage, heavy contamination
	B: Inadequate soft tissue wound and/or periosteal stripping
	C: open fracture with arterial injury and ischemia

Infection rate increases significantly as the grade of the injury increases (0-2% for type I, to 25-50% for type IIIC). As the grade increases, the pathogens shift towards nosocomial pathogens related to wound handling and colonization of open wounds by hospital bacterial flora. Thus, meticulous wound management greatly reduces infectious complications and has a greater impact than antibiotics delivered after the injury has occurred.

Protocol and Order Set:

Open Fractures:

1. Antibiotics:

- Types I and II:
 - a. Cefazolin 2 grams IV now and q8h x 3 doses
 - b. Penicillin allergic: Clindamycin 900 mg IV now and q8h x 3doses
- Type III (>10cm and/or severe contamination):
 - a. Vancomycin per EPIC dosing advisor x 24h
 - b. Ceftriaxone 2 grams IV now q24h x 1
 - c. Penicillin allergic: Levofloxacin 750mg IV now and q24h X 1
 - If multiple debridements are required, redose perioperatively with antibiotics utilized for initial Gustilo classification

****If open facial fracture, please refer to the Craniofacial PMG**

Ballistic Fractures:

1. Antibiotics:

- High Velocity: Treat as Grade III Gustilo (see regimens above)
- Low Velocity
 - a. Extremity (operative): Cefazolin 2 grams IV now and q8h x 3 doses perioperatively
 - b. Extremity (Non-operative): no prophylaxis required
 - c. Hip and pelvic fractures with concomitant bowel injury: Cefoxitin 2 grams IV now and q6h X 4 doses preoperatively
 - d. Hip and pelvic fractures with no bowel contamination:
 - Non-operative: no prophylaxis required
 - Operative: Cefazolin 2 grams IV now and q8h x 3 doses

References

1. Gosselin RA, Roberts I, Gillespie WJ. Antibiotics for preventing infection in open limb fractures. *Cochrane Database Syst Rev* 2004;CD003764.
2. Hauser CJ, Adams CA, Jr., Eachempati SR. Surgical Infection Society guideline: prophylactic antibiotic use in open fractures: an evidence-based guideline. *Surg Infect (Larchmt)* 2006; 7:379-405.
3. Luchette FA, Bone LB, Born CT, DeLong WG, Hoff WS, Mullins D, Palumbo F, Pasquale MD. EAST Practice Management Guidelines Work Group: Practice management guidelines for prophylactic antibiotic use in open fractures. Eastern Association for the Surgery of Trauma www.east.org, 1-28. 2000.
4. Dunkel N, Pittet D, Tovmirzaeva, et al. Short duration of antibiotic prophylaxis in open fractures does not enhance risk of subsequent infection. *Bone Joint J.* 2013;95-B:831-837.
5. Nguyen MP, Savakus JC, O'Donnell JA, et al. Infection rates and treatment of low-velocity extremity gunshot injuries. *J Orthop Trauma.* 2017;31:326-329.
6. Papasoulis E, Patzakis MJ, Zalavras CG. Antibiotics in the treatment of low-velocity gunshot-induced fractures. *Clin Orthop Relat Res.* 2013;471:3937-3944.
7. Vasanth S, Thakore RV, Stinner DJ, et al. Gunshot-induced fractures of the extremities: a review of antibiotic and debridement practices. *Curr Rev Musculoskelet Med.* 2015;8:276-289.

Revised December 2022
Jill Streams, MD
Robert Boyce, MD,
Leanne Atchison, PharmD
Jennifer Beavers, PharmD, BCPS

Approved: 1/25/23