

## CHEST PROTOCOL FOR BLUNT FORCE THORACIC TRAUMA

Managing patients with multiple rib fractures or flail chest requires significant health care resources including intensive care unit monitoring, ventilator management [1-3], surveillance and treatment of pneumonia, and pain management protocol. Both acute complications and long-term disabilities occur as a result of this very common pattern of thoracic trauma.

In a retrospective review of 64,750 civilian trauma patients in the National Trauma Databank (NTDB) with 1 or more rib fractures, mortality in the entire cohort was 10%[1]. Mortality increased stepwise with each additional fracture to a maximum of 34.4% with 8 or more fractures. Epidural analgesia was used in only 2.2% of patients but significantly reduced mortality in patients with 2 rib fractures and 4 or more rib fractures. For patients with 8 rib fractures, only 7% received an epidural catheter; however, those who did had a 98% survival as compared to a 65% survival without an epidural. Consideration for single or multiple paravertebral catheters should also be considered when an epidural catheter is contraindicated or technically unsuccessful as there is evidence of efficacy similar to epidurals with fewer contraindications and hypotension [4].

The nationwide opioid abuse epidemic has increased awareness of how acute pain is managed. A joint practice guideline from trauma surgeons and anesthesiologists conditionally recommends use of multimodal analgesia rather than opioids alone in patients with blunt thoracic trauma (10). Emphasis should be on reducing the duration and dosage of opioids as much as possible. Multimodal approaches involving non-steroidal anti-inflammatory drugs (NSAIDs) and GABA-analogs (gabapentin and pregabalin) have been shown to improve perioperative pain control and have been adapted as an initial approach for blunt chest wall injury [5-7].

Use of a rib fracture protocol has recently been shown to reduce ICU length of stay, hospital length of stay, and pneumonia in patients with rib fractures [3]. There was also a trend towards reduced mortality. This protocol involved a multidisciplinary approach involving the trauma team, anesthesia pain service, advanced care practitioners, physical therapy, and nutrition.

Long-term outcomes are similarly poor in these patients. Recent civilian studies indicate that the morbidity and lost productivity in patients suffering thoracic trauma is substantial even in the relatively young and those with minimal injuries. For example, patients with rib fractures are significantly more disabled at 30 days post injury when compared with patients with chronic medical illness and lose an average of 70 days of work or usual activity during their acute recovery [8].

From a functional standpoint, a recent study found that for up to 2 years after pulmonary contusion and rib fractures, multiple PFT parameters and exercise tolerance are compromised with VO<sub>2</sub>max of 60% predicted[9]. In patients with flail chest, 50 to 60% develop permanent morbidity with persistent chest wall pain or deformity being the most common long-term problems [10, 11]. As many as 40% of these patients still had not returned to work one year after their injury.

The challenge, therefore, is to identify treatment strategies that improve both short and long term outcomes in patients with severe chest wall trauma. We believe a multidisciplinary approach is warranted at VUH given the frequency of rib fractures in our patient population and the potential benefits to having a treatment protocol for managing acute fractures and a multi-modal approach to avoiding long-term disability (Appendix A).

Policy Statement:

1. This policy applies to trauma patients with rib fractures who are admitted to the Trauma service. Patients who meet the defined criteria will be managed according to this protocol as detailed in Appendix A.
2. The key pain management decisions are: 1) implementation of basic multimodal analgesia (acetaminophen, low-dose oral opioid, gabapentinoid, NSAID) unless contraindicated; 2) need for additional pain adjuncts beyond basic multimodal pain control (atypical oral medications, regional anesthesia techniques, IV analgesic infusions, etc.); 3) the need for operative stabilization of dislocated fractures or a flail chest; 4) patient suitability for regional anesthetic procedures; and 5) analgesic requirements at discharge (drug, dose, and duration). An attending trauma surgeon and attending anesthesiologist on the Acute Pain Service (APS) will be involved in these decisions and the APS plan will be communicated to the Trauma provider team.
3. If the patient has a flail chest, displaced rib fractures causing significant chest wall deformity, a need for surgical intervention in the chest in the setting of multiple rib fractures, or chronic pain from rib fractures, operative stabilization of the fractures should be considered. This does not replace the need for multimodal pain control.
4. Pain management beyond basic multimodal oral pain regimen will be managed primarily by APS. Management recommendations will be communicated to the Trauma team by the APS service. Regional anesthesia solutions and IV analgesic infusions (including but not limited to lidocaine and ketamine) will be ordered directly by APS. Management decisions such as placement or non-placement of regional anesthetics and the use of IV analgesic infusions will be communicated to the responsible trauma staff.


Appendix A: Vanderbilt Rib Fracture Protocol

VUMC Rib Fracture Management Pathway

- Supervised Incentive Spirometry (10 reps/hr, goal  $\geq$  15 mL/kg within best flow rate)
- RT evaluation
- PT consult for early mobility
- Maintain euvolemia

Criteria for Acute Pain Service (APS) consult:

- Age  $\geq$  45
- # of rib Fx  $>$  3
- Daily IS volume  $\leq$  15 mL/kg
- Pain (at rest and with cough)  $\geq$  5/10
- Present of admission medications:
  - $\geq$ 60 MME (milligram morphine equivalent ) per day
  - Current buprenorphine (Suboxone) or methadone use



- \*Multimodal Pain Therapy:
- NSAIDs
  - Tylenol
  - GABA-analogs (as indicated)
  - +/- opioids (oral preferred over IV; PCA preferred if IV required)

- RT evaluation includes assessment for:
- Adequacy of oxygenation/ventilation
  - Alveolar recruitment
  - Need for bronchodilators
  - Secretion management

## Appendix B: APS Consult Guidelines

### APS Consult considerations

#### The APS will:

- Evaluate patients within 6 hours of consultation
- Reassess 18-36 hours after initial evaluation
- Leave epidural or other perineural catheter in place for 3-5 days, up to 7 for special circumstances
- Provide daily follow-up while TEC in place or IV infusions running
- Consider alternative blocks if epidural contraindicated or unilateral injury
- Consider analgesic infusion if anesthetic block contraindicated
- Consider for intubated patients (RASS -2 to +1) with anticipated extubation in the next 24hrs
- Discuss with Trauma Faculty if regional anesthesia contraindicated

### Thoracic Epidural Contraindications

- Labs: Platelets < 80,000, INR >1.5, elevated PTT
- Medications: anticoagulants, antiplatelet agents, or the inability to rule out the use of these medications (see Appendix C)
- Infection: Systemic or insertion site infection
- Certain traumatic injuries: Epidural or spinal cord hematoma, TBI with midline shift, spinal cord injury, operative spine fractures adjacent to insertion site
- Epidural contraindications may not preclude other blocks

### Barriers to Regional Anesthesia – though not contraindications, these may make regional anesthesia challenging or require changes in timing or technique

- Deep sedation: < RASS -3
- Injuries requiring placement in traction (acetabular fracture, etc.)
- Patient distant from extubation or ventilator weaning
- Inability to provide consent or identify surrogate decision-maker
- Operative spine fractures (away from insertion site) or pending spine evaluation

## Appendix C: Anticoagulation guidelines for regional anesthesia

American Society of Regional Anesthesia and Pain Medicine (ASRA) Guidelines  
<https://www.asra.com/advisory-guidelines/article/9/regional-anesthesia-in-the-patient-receiving-antithrombotic-or-thrombolytic-ther>

Medication	Prior to procedure	With catheter in place	Prior to removal	After catheter removal
Heparin				
SC Heparin TID ( $\leq 5000$ units)	Hold AM dose	Hold until 1 hour after placement	Hold AM dose	Resume 1 hour after removal
SC Heparin ( $>5000$ units)	Hold AM dose, wait until PTT $<40$	AVOID WHILE CATHETER IN PLACE		
IV Heparin	Hold infusion 4-6 hrs, wait until PTT $<40$			
Low Molecular Weight Heparin				
Prophylactic dose (30 mg SQ BID, 40 mg SQ BID)	Hold 12 hrs	AVOID WHILE CATHETER IN PLACE		Resume 4 hours after removal
Therapeutic dose (1 mg/kg SC BID or 1.5 mg/kg daily)	Hold 24 hrs			
Vitamin K antagonist				
Warfarin	Hold until INR $<1.5$	AVOID WHILE CATHETER IN PLACE	INR $<1.5$	May start immediately
Anti-platelets				
Aspirin	NO RESTRICTIONS			
Plavix	Hold for 7 days	AVOID WHILE CATHETER IN PLACE		Hold loading dose for 6 hours after removal. If NO LOADING DOSE, may start immediately.

References:

1. Fligel BT, Luchette FA, Reed RL, Esposito TJ, Davis KA, Santaniello JM, et al. Half-a-dozen ribs: the breakpoint for mortality. *Surgery*. 2005;138(4):717-23; discussion 23-5. Epub 2005/11/05. doi: S0039-6060(05)00433-2 [pii]  
10.1016/j.surg.2005.07.022. PubMed PMID: 16269301.
2. Sharma OP, Oswanski MF, Jolly S, Lauer SK, Dressel R, Stombaugh HA. Perils of rib fractures. *Am Surg*. 2008;74(4):310-4. Epub 2008/05/06. PubMed PMID: 18453294.
3. Todd SR, McNally MM, Holcomb JB, Kozar RA, Kao LS, Gonzalez EA, et al. A multidisciplinary clinical pathway decreases rib fracture-associated infectious morbidity and mortality in high-risk trauma patients. *Am J Surg*. 2006;192(6):806-11. Epub 2006/12/13. doi: S0002-9610(06)00602-7 [pii]  
10.1016/j.amjsurg.2006.08.048. PubMed PMID: 17161098.
4. Mohta M, Verma P, Saxena AK, Sethi AK, Tyagi A, Girotra G. Prospective, randomized comparison of continuous thoracic epidural and thoracic paravertebral infusion in patients with unilateral multiple fractured ribs--a pilot study. *J Trauma*. 2009;66(4):1096-101. Epub 2009/04/11. doi: 10.1097/TA.0b013e318166d76d. PubMed PMID: 19359920.
5. Chou R, Gordon DB, de Leon-Casasola OA, Rosenberg JM, Bickler S, Brennan T, et al. Management of Postoperative Pain: A Clinical Practice Guideline From the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *J Pain*. 2016;17(2):131-57. Epub 2016/02/02. doi: 10.1016/j.jpain.2015.12.008. PubMed PMID: 26827847.
6. Galvagno SM, Jr., Smith CE, Varon AJ, Hasenboehler EA, Sultan S, Shaefer G, et al. Pain management for blunt thoracic trauma: A joint practice management guideline from the Eastern Association for the Surgery of Trauma and Trauma Anesthesiology Society. *J Trauma Acute Care Surg*. 2016;81(5):936-51. Epub 2016/10/22. doi: 10.1097/TA.0000000000001209. PubMed PMID: 27533913.
7. Dennis BM, Bellister SA, Guillaumondegui OD. Thoracic Trauma. *Surg Clin North Am*. 2017;97(5):1047-64. Epub 2017/09/30. doi: 10.1016/j.suc.2017.06.009. PubMed PMID: 28958357.
8. Kerr-Valentic MA, Arthur M, Mullins RJ, Pearson TE, Mayberry JC. Rib fracture pain and disability: can we do better? *J Trauma*. 2003;54(6):1058-63; discussion 63-4. Epub 2003/06/19. doi: 10.1097/01.TA.0000060262.76267.EF. PubMed PMID: 12813323.
9. Amital A, Shitrit D, Fox BD, Raviv Y, Fuks L, Terner I, et al. Long-term pulmonary function after recovery from pulmonary contusion due to blunt chest trauma. *Isr Med Assoc J*. 2009;11(11):673-6. Epub 2010/01/30. PubMed PMID: 20108554.
10. Beal SL, Oreskovich MR. Long-term disability associated with flail chest injury. *Am J Surg*. 1985;150(3):324-6. Epub 1985/09/01. PubMed PMID: 4037191.
11. Landercasper J, Cogbill TH, Lindesmith LA. Long-term disability after flail chest injury. *J Trauma*. 1984;24(5):410-4. Epub 1984/05/01. PubMed PMID: 6716518.