

Practice Management Guideline for Speech Therapy/Cog Evaluations

Purpose: To define patient populations which necessitate a cognitive evaluation in the setting of a Traumatic Brain Injury

Indications:

- Anyone with TBI (minimal, mild, moderate, severe) should have a cognitive evaluation ordered.
- Positive LOC alone should not trigger an order for a cognitive evaluation.

If the patient is medically ready for discharge, and the only barrier to discharge is a cognitive evaluation, the following inclusion/exclusion criteria should be utilized:

A cognitive evaluation may be deferred if:

1. the patient has baseline dementia requiring supervision (excluded if they live independently)
2. GCS 15 upon arrival and throughout hospitalization (regardless of LOC status)

A cognitive evaluation should be done if:

1. the patient is enrolled in any type of school (high school, college, ect)
2. the patient is exhibiting any concussive symptoms: tinnitus, HA, diplopia, N/V, repetitive questioning.

If a patient with TBI is discharged prior to obtaining a cognitive evaluation:

1. Schedule them in trauma APP clinic in 14 days and TBI clinic in 3 months
2. Give patient TBI outpatient resources handout

Authors:

Katie Forrester, ACNP

Amanda Hereford, MA, CCC-SLP

Melissa Morton, ACNP

Manry Martin, ACNP

Brad Dennis, MD

Jill Streams, MD

Revision Dates:

8/2023