

Trauma ICU Nutrition Management Guidelines

Trauma Critical Care Nutrition Guidelines

Clinical judgment may supersede guidelines as patient circumstances warrant

ASSESSMENT AND EVALUATION

- All patients admitted to the Trauma Intensive Care Unit require a nutrition risk assessment within 24 hours and a nutrition plan within 48 hours
- Consult Nutrition Service as needed for specific recommendations (i.e., tube feeding formulations, oral supplements, poor oral intake, education)

ADMINISTRATION

- Enteral nutrition (EN) preferred over parenteral nutrition (PN)
- Reduce risk of aspiration by reducing sedation, elevating HOB 30 – 45 degrees, performing mouth care per VAP Guidelines and minimizing transport out of ICU

Oral Nutrition

- Oral intake preferred method of nutrition if appropriate for patient
- Initiate regular diet with oral diet advancement with addition of oral supplement to optimize po intakes

Enteral Nutrition

- Initiate EN 24 – 48 hours following onset of critical illness and admission to ICU, after resuscitation efforts completed and/or hemodynamic stability achieved
- Initiate tube feedings and advance as quickly as tolerated in 24 – 48 hours to goal within 48 – 72 hours
 - Weaning EN (transitioning to PO diet)
 - Cycle EN x 12hr, 7p to 7am (for 50% of needs during first few days of transition)
 - Wean off EN once patient consistently consumes and tolerates on average 50% or more of meals
- Post Pyloric access preferable if EN access needed, especially with high aspiration risk, but nutrition should not be delayed if only gastric access obtained.
 - Access
 - Gastric
 - Short term: Orogastric tube (OGT), Nasogastric tube (NGT), Dobhoff tube (DHT)
 - Long term: Percutaneous endoscopic gastrostomy (PEG) or laparoscopic gastrostomy
 - Post-pyloric:
 - Short term: DHT (confirmed by abdominal radiographic imaging (KUB))
 - Long term: PEG-Jejunostomy (for unsuccessful placement DHT for post-pyloric access)

Parenteral Nutrition

- If low nutrition risk and unable to meet > 60% energy and protein requirements via EN within 7 - 10 days, then initiate PN
- If high nutrition risk present (malnutrition upon admission, inability to use GI tract expected for more than 3-5 days) and EN not feasible, initiate PN as soon as possible after resuscitation efforts completed.
- If high nutrition risk present (malnutrition upon admission determined by AND/ASPEN criteria and inability to use GI tract expected for more than 3-5 days), initiate PN as soon as possible after resuscitation efforts completed
- Wean TPN when 60% of TF goal met or 60% of meals consumed
 - Decrease TPN to ~half, decrease dextrose/AA per PN team order
 - Wean off TPN as TF rate advances or per clinical judgment

Revisions: 4/20/2021, 4/2023

Authors:

Beth Mills, MS, RD, CNSC, LDN
Laurie Ford, APNP-BC
Stephen Gondek, MD

If LOS>7days and pt has not consistently met on average near 60% estimated needs, consider nutritional provision from a combination of PO/EN/PN routes.

DOSING

- Dosing weight
 - Use actual weight or known usual body weight (UBW) if BMI <29.99 for height.
 - Use upper ideal body weight (IBW) for height if BMI 30 or greater.
 - Hamwi Method
 - Men: 106# (48kg) for 1st 5 feet, then add 6# (2.7kg) per inch >5 feet, +/-10%
 - Women: 100# (45kg) 1st 5 feet, then add 5# (2.3kg) per inch >5feet, +/-10%
 - Use actual body weight if weight < IBW
- Energy goals:
 - 25 kcal/kg dosing weight/day
 - If BMI >30 (Class I, Class II or Class III Obesity), use 25 kcal/kg upper IBW/day
- Protein goals:
 - General 1.2 – 2.0 g/kg dosing weight/day
 - Obesity
 - If BMI 30 –39.99, use 2g/kg upper IBW/day
 - If BMI > 40, use 2.5g/kg upper IBW/day
 - Renal Failure:
 - HD 1.5 - 2.0 g/kg dosing weight
 - CRRT: 1.5 -2.0 with maximum 2.5g/kg dosing weight
 - Hepatic Failure: 1.2 - 2.0/kg dry or actual body weight/day
 - Spinal Cord Injury: 2.0/kg dosing weight
 - Traumatic Brain Injury: 1.5-2.0/kg dosing weight
 - Open Abdomen with negative pressure therapy estimated protein loss: add 2.9gm/liter of exudate lost

MONITORING

- Serum protein markers (i.e. prealbumin, CRP) not recommended for evaluation of nutritional status or goals
- GI Intolerance
 - Gastric residual volume (GRV) not utilized as routine evaluation of tolerance. Daily physical examination, patient symptoms, clinical risk factors, and abdominal radiographic films should be utilized to determine tolerance. If checked, Tube feeds should not be held unless GRV is > 500 cc
 - Prokinetic agents may be introduced if GI intolerance suspected or for patients with high risk of aspiration. Consider QTc prolongation.
 - Erythromycin 200mg IV or per tube q6h x 3 days
 - Metoclopramide 10mg IV q6h x 3 days
 - Naloxone 8mg q8h x 3 days, then 8mg q6h prn
- For persistent diarrhea and C. Diff infection ruled out, initiate Nutrisource fiber 4 packets in 24 hours
- Special considerations
 - **Refeeding syndrome**
 - Replete electrolytes, provide thiamine, folic acid and MVI prior to initiation of tube feedings
 - Patients at risk for refeeding syndrome, initiate trophic feedings (no more than 25% of goal) and then check BMP, phosphorus and magnesium levels
 - Advance tube feedings slowly over 3 – 4 days
 - Check BMP, phosphorus and magnesium levels daily as EN advances to goal
 - **Open Abdomen**
 - Early EN recommended 24 – 48 hours after injury, without evidence of bowel injury

Hyperglycemia: (VUMC EN formulary does not have a “diabetic” EN formula) per gram protein provided

Revisions: 4/20/2021, 4/2023

Authors:

Beth Mills, MS, RD, CNSC, LDN
Laurie Ford, APNP-BC
Stephen Gondek, MD

Impact Peptide 1.5 or Peptamen Intense VHP will provide lowest amount of carbohydrate per TF goal.

ASSOCIATED MDSCC PROTOCOLS

- Glycemic Protocol
- Gastrointestinal Stress Ulcer Prophylaxis
- VAP Protocol

Appendix 1


TICU ENTERAL NUTRITION TUBE FEEDING FORMULATIONS

<p>Critically Ill Patient</p> <p>Impact Peptide 1.5 (high protein as peptides, immune enhancing formula)</p>	<p>Obese Critically Ill Patient</p> <p>Peptamen Intense VHP (very high protein formula)</p>	<p>Non-Critically Ill Patient</p> <p>Isosource HN Nutren 1.5 Nutren 2.0</p>
---	--	--

Consult Nutrition Service for disease specific formulations in TICU

<p>Respiratory failure</p> <p>Nutren 2.0 (Volume restricted formula)</p>	<p>Admitted with pre-existing renal failure</p> <p>Novasource Renal (Electrolyte restricted formula)</p>	<p>Renal failure Develops</p> <p>HD: Novasource Renal</p> <p>CRRT: Impact Peptide 1.5 or Peptamen Intense VHP</p>	<p>Acute Pancreatitis</p> <p>Peptamen 1.5</p>	<p>MODS/Chyle Leak</p> <p>Vivonex RTF 1.0</p>	<p>Modulars</p> <p>Nutrisource Fiber</p> <p>Prostat Max (Protein)</p>
---	--	--	--	--	--

INTERNAL/PARENTERAL NUTRITION FLOW DIAGRAM



PREOPERATIVE ENTERAL NUTRITION PROTOCOL FOR PATIENTS WITH PROTECTED AIRWAY (Trach/Oral ETT)

NON-ABDOMINAL SURGERY

- Turn tube feedings off just prior to OR departure or bedside procedure
- Gastric tube will be flushed and aspirated

ABDOMINAL SURGERY OR OPERATIVE INTERVENTION REQUIRING PRONE POSITIONING

- Turn tube feedings off 6 hours before planned anesthesia
- Gastric tube will be flushed and aspiration prior to OR departure

UPPER GI ENDOSCOPY

- Turn tube feedings off 1 hour prior to elective endoscopy
- Place NGT to suction

OTHER CONSIDERATIONS

- Stop insulin infusion prior to OR transport
- Alert anesthesiology to perform accucheck perioperatively in OR if SQ insulin given within 2 hours
- Restart tube feedings post-surgery unless orders to hold post-surgery
- Patient with confirmed post-pyloric feeding tube, consider perioperative continuous feeding by anesthesiology and surgeon

Sources for Guideline Development:

- Boullata JI, Carrera AL, Harvey LH, Hudson L, et al. ASPEN Safe Practices for Enteral Nutrition Therapy. *Journal of Parenteral and Enteral Nutrition*. 2017; 41(1):15 - 103.
- McClave SA, Taylor, BE, Martindale RG, Warren MM, et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society of Parenteral and Enteral Nutrition (ASPEN). *Journal of Parenteral and Enteral Nutrition*. 2016; 40 (2): 159-211.
- Taylor BE, et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (ASPEN). *Critical Care Medicine*. 44(2): 390 – 438, February 2016.
- Kohn JB. Adjusted or Ideal Body Weight for Nutrition Assessment? <http://dx.doi.org/10.1016/j.jand.2015.02.007>
- Andrews, AM, Pruziner, AL. Guidelines for Using Adjusted versus Unadjusted Body Weights When Conducting Clinical Evaluations and Making Clinical Recommendations. <http://dx.doi.org/10.1016/j.jand.2016.07.003>
- Wade C, Wolf, SE, Reuben S, et al. Loss of Protein, immunoglobulins, and Electrolytes in Exudates From Negative Pressure Wound Therapy. *Nutrition in Clinical Practice* 2010;25 (5):510-516.
- Compher C, Bingham AL, McCall M, et al. Guideline for the provision of nutrition support therapy in adult critically ill patient: The American Society for Parenteral and Enteral Nutrition. *Journal of Parenteral and Enteral Nutrition*. 2022; 46:12-41. DOI:10.1002/jpen.2267
- Schwartz DB, Barrocas A, Annetta MG, et al. Ethical Aspects of Artificially Administered Nutrition and Hydration: An ASPEN Position Paper. *Journal of Parenteral and Enteral Nutrition*. 2021;35 (2): 254-267. DOI: 10.1002/ncp.10633
- Bechtold, ML, Brown PM, Escuro A, et al. When is enteral nutrition indicated? *Journal of Parenteral and Enteral Nutrition*. 2022; 46:1470-1496. DOI:10.1002/jpen.2364
- Singer P, Blaser AR, Berger MM, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clinical Nutrition*. 2019;38:48-79. <https://doi.org/10.1016/j.clnu.2018.08.037>
- Academy of Nutrition and Dietetics. Adult Nutrition Care Manual, Nutrition Care, Critical Illness 2021 update. <http://www.nutritioncaremanual.org>. accessed 9/10/22