



TRAUMA GLYCEMIC CONTROL PROTOCOL

Initial Management: Triage of Care and Initial Treatment



Euglycemic Management: Insulin



DIVISION OF ACUTE CARE SURGERY

Calculating Basal and Prandial Insulin Regimen:

- Basal Dose Adjustment: $Basal\ Dose = \frac{\text{Total insulin requirement over prior 24h}}{2}$
- Receiving continuous tube feeds (in TICU): $Prandial\ Dose = \frac{\text{Total insulin requirement over prior 24h}}{2} \div 6$
- Receiving continuous tube feeds (on stepdown): $Prandial\ Dose = \frac{\text{Total insulin requirement over prior 24h}}{2} \div 4$
- Receiving carbohydrate-controlled diet: $Prandial\ Dose = \frac{\text{Total insulin requirement over prior 24h}}{2} \div 3$
- Ensure trauma SSI is ordered as /15 or /12 if adding prandial coverage.

Important Considerations:

- If NPO or tube feedings held, hold prandial insulin but continue basal and sliding scale insulin
- All patients with insulin orders need the hypoglycemia protocol off the order set
- SSI alone without basal insulin is not recommended for Type 1 and Type 2 Diabetes
 - USE CAUTION in patients with renal dysfunction, elderly, and those with large volume fluid shifts.
Recommend using lower insulin doses
- Consider discontinuing blood glucose monitoring and sliding scale insulin if BG remains ≤ 150 mg/dL AND tube feed goals met (or patient tolerating a diet) for 24 hours and off vasopressors

References:

1. Yendamuri, Saikrishma, et al. Admission hyperglycemia as a prognostic indicator in trauma. J Trauma 2003; 55:33-38.
2. Laird A, et al. Relationship of early hyperglycemia to mortality in trauma patients. J Trauma 2004; 56:1058-1062.
3. Sung J, et al. Admission hyperglycemia is predictive of outcome in critically ill trauma patients. J Trauma 2005; 59:80-83.
4. Van den Berghe G, Wouters P, Weekers F, et al. Intensive Insulin Therapy in Critically Ill Patients. New England Journal of Medicine 2001; 345:1359–1367.
5. The NICE-SUGAR Study Investigators, Finfer S, Chittock DR, et al. Intensive versus Conventional Glucose Control in Critically Ill Patients. New England Journal of Medicine 2009; 360:1283–1297.
6. Jacobi J, Bircher N, Krinsley J, et al. Guidelines for the use of an insulin infusion for the management of hyperglycemia in critically ill patients. Critical Care Medicine 2012; 40:3251–3276.
7. Mowery NT, Guillaumondegui OD, Gunter OL, et al. Severe hypoglycemia while on intensive insulin therapy is not an independent predictor of death after trauma. The Journal of Trauma: Injury, Infection, and Critical Care 2010; 68:342–347.
8. Mowery NT, Gunter OL, Kauffmann RM, et al. Duration of time on intensive insulin therapy predicts severe hypoglycemia in the surgically critically ill population. World Journal of Surgery 2011; 36:270–277.
9. Krinsley J, et al. Glycemic variability: A strong predictor of mortality in critically ill patients. Crit Care Med 2008; 36:3008-3013.
10. Kauffmann RM, Hayes RM, Jenkins JM, et al. Provision of Balanced Nutrition Protects Against Hypoglycemia in the Critically Ill Surgical Patient. Journal of Parenteral and Enteral Nutrition 2011; 35:686–694

DIVISION OF ACUTE CARE SURGERY

11. Bode, BW, Braithwaite SS, Steed RD, Davidson PC. Intravenous insulin infusion therapy: indications, methods, and transition to subcutaneous insulin therapy. *Endocrine Practice* 2004; 10(2):71-80.

Last revised: October 2025, March 2023, January 2020

Revision Team:

Jill Streams, MD- Trauma PI Director

Bradley Dennis, MD- Trauma Medical Director

Caroline Banes, DNP, ACNP-BC

Kayla Harding, MSN, AGACNP

Jennifer Beavers, PharmD

Chelsea Taska, PharmD