

**VANDERBILT UNIVERSITY MEDICAL CENTER  
DIVISION OF TRAUMA AND SURGICAL CRITICAL CARE**

**Bedside Surgery Protocol**

---

---

1. Indications
  - a. Decompressive celiotomy for abdominal compartment syndrome
  - b. Exploratory celiotomy for acute hemodynamic decompensation due to hemorrhage
    - i. should be reserved for patients who are prohibitive risks for OR transport
  - c. Re-exploration of a previous open abdomen for dressing change or closure
  - d. Exploratory celiotomy to rule out intra-abdominal sepsis in a patient whose physiologic condition prohibits safe transport to the operating room
  - e. Percutaneous tracheostomy
  - f. Percutaneous gastrostomy
  - g. Bronchoscopy
  - h. Decompressive colonoscopy
2. Bedside surgery protocol: General
  - a. Critical care attending and the operating surgeon should be present for the entire surgical procedure
  - b. Critical care attending will oversee the anesthetic management of the patient
    - i. intravenous general anesthesia
      - a. anxiolysis
        - i. e.g. propofol, Versed, Ativan, ketamine
      - b. analgesia
        - i. e.g. fentanyl, morphine, hydromorphone, ketamine
      - c. chemical paralysis as needed
        - i. vecuronium, cisatracurium, rocuronium
    - ii. optimize ventilator settings
      1. place patient on mandatory rate at 100% FiO<sub>2</sub>
      2. volume control preferred, particularly for percutaneous tracheostomy
  - c. Obtained informed consent if procedure not emergent
  - d. Pre-procedure timeout to be performed by surgical team, procedure support staff, and bedside nursing
    - i. surgical team availability
    - ii. laterality or level
    - iii. verified patient medical record number
    - iv. review allergies
    - v. procedure verification
    - vi. SCIP criteria?
    - vii. informed consent
    - viii. appropriate instruments available
    - ix. special considerations
  - e. Bedside nurse and respiratory therapist will monitor the patient and record the procedure vital signs on conscious sedation sheet
    - i. monitors: ECG, blood pressure (arterial line/Q5 minutes), pulse oximetry, ICP as indicated, ventilator settings
  - f. Critical care or procedure attending signs the sedation sheet post-procedure
  - g. Sterile perimeter will be set up in the patient's room.

- h. All individuals within sterile perimeter must wear personal protective equipment
    - i. i.e. surgical cap, mask, eye protection
  - i. All members of the operative team must decontaminate hands as per OR routine
  - j. prophylactic antibiotics indicated only if new surgical wound is made
    - i. antibiotic choice per surgeon preference based on degree of case contamination
  - k. Chlorprep agent of choice for skin preparation unless contraindicated
  - l. Indication to proceed to the operating room (level 1)
    - i. surgeon preference
    - ii. uncontrollable hemorrhage
    - iii. instrumentation requirement exceeding bedside capability
  - m. Procedures are documented in Star forms by participating house staff and signed by Attending staff in a timely fashion.
1. Bedside laparotomy considerations
    - a. Electrocautery will be available as needed.
    - b. Wall suction canisters available with tubing and Yankauer tips.
    - c. 4 L of warm crystalloid solution available.
    - d. A standard bedside celiotomy tray including suture will be set up on the sterile field.
    - e. Vacuum pack changes for damage control celiotomy
      - i. every 48-72 hours
      - ii. typical dressing includes bowel isolation bag, safety towels with radiographic marker, 10/19 Fr. JP drains, and adhesive barrier dressing
      - iii. proprietary dressings may be used as suitable (KCI Abthera)
  2. Bedside tracheostomy considerations
    - a. High risk patients must be identified preprocedure.
      - i. Morbid obesity
      - ii. airway edema
      - iii. cervical trauma
      - iv. extremes of age
      - v. other considerations
        1. Mandibulomaxillary fixation
        2. Halo brace
        3. High ventilator settings
          - a. FIO<sub>2</sub> >50%
          - b. PEEP >10
        4. bleeding diathesis
        5. anatomical considerations
    - b. Consideration should be given to bronchoscopic guidance versus conversion to open procedure either a bedside or in the operating room
    - c. Proximal XLT tracheostomy should be selected for patients with significant obesity
    - d. Blue Rhino percutaneous tracheostomy kit, cutdown instrument set, appropriate suture, tracheostomy soft pack, tracheostomy tube (typically 8 Shiley), intubation tray, and tidal CO<sub>2</sub> monitor, difficult airway bag.
    - e. Withdrawal of endotracheal tube must be performed by experienced personnel with care to avoid inadvertent extubation.
    - f. Confirmation of endotracheal placement of tracheostomy is by physical examination, CO<sub>2</sub> color change, and inspired tidal volume ~expired tidal volume.

- g. Tracheostomy secured with suture and neck strap.
  - h. Post procedure chest x-ray is recommended.
3. Bedside PEG considerations
- a. Video gastroscopy.
  - b. Bard PEG (pull-type) feeding tube kit, 20-French.
  - c. T-fasteners may be utilized for patients with malnutrition or immunosuppression.
  - d. T-fasteners; these are to be cut no later than date 10.
  - e. Cutdown instrument set and appropriate suture (e.g. 2-0 nylon or silk).
  - f. Notation is made regarding site of gastrostomy tube as well as depth both in the procedure as well as the nursing note.

*Revised 2011, Oliver Gunter, M.D., FACS*

*Reviewed December 2013:*

*Oliver Gunter, MD, FACS*

*Laurie Ford ACNP-BC*

*Caroline Banes ACNP-BC*

*Approved:*

---

*Addison May, MD*