Vanderbilt University Medical Center Multidisciplinary Surgical Critical Care

Standard Operating Procedure: Respiratory Therapy / Critical Care Communication

Purpose:

To standardize the process of ventilator order entry and communication with respiratory therapist to facilitate consistent and uniform ventilatory strategy for critically ill patients in the SICU

Procedure:

- 1. Print out a **Respiratory Therapy List** from WIZ prior to rounds.
 - In Wiz Order, go to the **Printouts** and select **Respiratory Therapy List**, then select (SICU) **scratch census**, select **all**, then select **OK** and print.
- 2. Review the respiratory care orders as printed on the **Respiratory Therapy List** for the SICU consult scratch census on each patient during rounds.
- **3.** Update orders as needed using the rolling computer while rounding.
 - For ventilator weaning, three (3) order sets are present in WIZ and can be accessed by typing in vent wean and then selecting SICU ventilator weaning order set.
 - o Choose the appropriate wean: **PSV Wean, FiO2 Wean** or **PEEP Wean.**
 - o These orders can be modified once these orders have been entered
 - Additional orders that require entry are:
 - o Trach collar trials with frequency and duration,
 - o extubation to a desired FiO2
 - o incentive spirometry with frequency.
- 4. Print out an updated **Respiratory Therapy List** at the completion of rounds, senior member of MDSCC team to discuss plans of care with the therapist.
 - SICU respiratory therapist pager 835-5990 to round and go over the plan of care for each patient.
 - Please be aware without this direct communication, a delay may occur in therapists viewing of orders

Approved:	Revised : 3-1-2004
Addison K. May, MD, FACS	
Medical Director, SICU	